

Leukaemia Section

Short Communication

t(8;14)(q11;q32)

Jean-Loup Huret

Genetics, Dept Medical Information, University of Poitiers, CHU Poitiers Hospital, F-86021 Poitiers, France (JLH)

Published in Atlas Database: November 1999

Online updated version : <http://AtlasGeneticsOncology.org/Anomalies/t0814ID1112.html>

DOI: 10.4267/2042/37562

This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 2.0 France Licence.

© 1999 *Atlas of Genetics and Cytogenetics in Oncology and Haematology*

Clinics and pathology

Disease

Acute lymphoblastic leukemia (ALL) most often (14 cases); chronic myelogenous leukemia (CML) (3 cases); one case of histiocyte-rich B-cell lymphoma.

Etiology

Strikingly, of 18 patients, 4 have Down syndrome, 1 has neurofibromatosis Type I, and another one is dysmorphic and mentally retarded.

Epidemiology

Highly unbalanced sex ratio (13M/2F).

Clinics

Still poorly known.

Cytogenetics

Cytogenetics morphological

Sole anomaly in 4 ALL cases; accompany a t(9;22)(q34;q11) in 4 of the 14 ALL cases (and in the CML cases); unbalanced form with a der(14) t(8;14) in 3 cases, indicating that the crucial event is likely to lie on der(14).

Additional anomalies

t(8;14) seems to be typically an anomaly secondary to t(9;22) (7/18 cases (40%), see above); anomalies additional to t(8;14) are +X, and +8 (2 cases each).

Genes involved and proteins

Note

The gene involved in 8q11 is unknown; the gene involved in 14q32 is IgH, found rearranged in a case where it was tested.

References

Hayata I, Sakurai M, Kakati S, Sandberg AA. Chromosomes and causation of human cancer and leukemia. XVI. Banding studies of chronic myelocytic leukemia, including five unusual Ph11 translocations. *Cancer*. 1975 Oct;36(4):1177-91

Kardon NB, Slepowitz G, Kochen JA. Childhood acute lymphoblastic leukemia associated with an unusual 8;14 translocation. *Cancer Genet Cytogenet*. 1982 Aug;6(4):339-43

Carroll AJ, Castleberry RP, Crist WM. Lack of association between abnormalities of the chromosome 9 short arm and either "lymphomatous" features or T cell phenotype in childhood acute lymphocytic leukemia. *Blood*. 1987 Mar;69(3):735-8

Crist W, Carroll A, Shuster J, Jackson J, Head D, Borowitz M, Behm F, Link M, Steuber P, Ragab A. Philadelphia chromosome positive childhood acute lymphoblastic leukemia: clinical and cytogenetic characteristics and treatment outcome. A Pediatric Oncology Group study. *Blood*. 1990 Aug 1;76(3):489-94

Hayashi Y, Pui CH, Behm FG, Fuchs AH, Raimondi SC, Kitchingman GR, Mirro J Jr, Williams DL. 14q32 translocations are associated with mixed-lineage expression in childhood acute leukemia. *Blood*. 1990 Jul 1;76(1):150-6

Hu N, Bian ML, Le Beau MM, Rowley JD. Cytogenetic analysis of 51 patients with chronic myeloid leukemia. *Chin Med J (Engl)*. 1990 Oct;103(10):831-9

Wodzinski MA, Watmore AE, Lilleyman JS, Potter AM. Chromosomes in childhood acute lymphoblastic leukaemia: karyotypic patterns in disease subtypes. *J Clin Pathol*. 1991 Jan;44(1):48-51

Pui CH, Carroll AJ, Raimondi SC, Schell MJ, Head DR, Shuster JJ, Crist WM, Borowitz MJ, Link MP, Behm FG. Isochromosomes in childhood acute lymphoblastic leukemia: a collaborative study of 83 cases. *Blood*. 1992 May 1;79(9):2384-91

Secker-Walker LM, Hawkins JM, Prentice HG, Mackie PH, Heerema NA, Provisor AJ. Two Down syndrome patients with an acquired translocation, t(8;14)(q11;q32), in early B-lineage acute lymphoblastic leukemia. *Cancer Genet Cytogenet*. 1993 Oct 15;70(2):148-50

Testoni N, Zaccaria A, Martinelli G, Pelliconi S, Buzzi M, Farabegoli P, Panzica G, Tura S. t(8;14)(q11;q32) in acute lymphoid leukemia: description of two cases. *Cancer Genet Cytogenet*. 1993 May;67(1):55-8

Shearer P, Parham D, Kovnar E, Kun L, Rao B, Lobe T, Pratt C. Neurofibromatosis type I and malignancy: review of 32 pediatric cases treated at a single institution. *Med Pediatr Oncol*. 1994;22(2):78-83

Litz CE, Davies S, Brunning RD, Kueck B, Parkin JL, Gajl Peczalska K, Arthur DC. Acute leukemia and the transient myeloproliferative disorder associated with Down syndrome: morphologic, immunophenotypic and cytogenetic manifestations. *Leukemia*. 1995 Sep;9(9):1432-9

Schoch C, Rieder H, Stollmann-Gibbels B, Freund M, Tischler HJ, Silling-Engelhardt G, Fonatsch C. 17p anomalies in lymphoid malignancies: diagnostic and prognostic implications. *Leuk Lymphoma*. 1995 Apr;17(3-4):271-9

Lee AC, Chan LC, Kwong KW. Down syndrome, acute lymphoblastic leukemia, and t(8;14)(q11;q32) *Cancer Genet Cytogenet*. 1996 May;88(1):92

Sun T, Susin M, Tomao FA, Brody J, Koduru P, Hajdu SI. Histiocyte-rich B-cell lymphoma. *Hum Pathol*. 1997 Nov;28(11):1321-4

Forrest DL, Nevill TJ, Horsman DE, Brockington DA, Fung HC, Toze CL, Conneally EA, Hogge DE, Sutherland HJ, Nantel SH, Shepherd JD, Barnett MJ. Bone marrow transplantation for adults with acute leukaemia and 11q23 chromosomal abnormalities. *Br J Haematol*. 1998 Dec;103(3):630-8

Whitehead VM, Vuchich MJ, Cooley LD, Lauer SJ, Mahoney DH, Shuster JJ, Payment C, Koch PA, Akabutu JJ, Bowen T, Kamen BA, Ravindranath Y, Emami A, Look AT, Beardsley GP, Pullen DJ, Camitta B. Accumulation of methotrexate polyglutamates, ploidy and trisomies of both chromosomes 4 and 10 in lymphoblasts from children with B-progenitor cell acute lymphoblastic leukemia: a Pediatric Oncology Group Study. *Leuk Lymphoma*. 1998 Nov;31(5-6):507-19

This article should be referenced as such:

Huret JL. t(8;14)(q11;q32). *Atlas Genet Cytogenet Oncol Haematol*. 1999; 3(4):196-197.
